


Isolation shape (pt and eta dependence)

Y.S. Chung, Jiyeon Han, Arie Bodek
(University of Rochester)
Feb, 18, 2010 / LPC muons

Isolation Shape


- Pt and Eta dependence
 - Pooja checked isolation profile in Pt and Eta



- We need to study these dependence
 - check isolation profile in $\text{isoE} < 2.0$
 - check IsoR03 (LPC standard) and IsoR03fw (CMS standard)

IsoR03 Profile in Pt and Eta (W boson)



- Isolation profile in Pt and Eta (iso<2.0) - IsoR03



- Pt dependence is very small
- Eta dependence is large and similar structure with Pooja's plot
- IsoR03 is asymmetric between $\eta > 1.3$ and $\eta < -1.3$

IsoR03fw Profile in Pt and Eta (W boson)

- Isolation profile in Pt and Eta ($\text{iso} < 2.0$) - IsoR03fw




- Iso profile in Pt is relatively flat
- No eta dependence in the endcap region
- IsoR03fw is better to use considering eta dependence
 - need to check what correction makes big eta dependence in isoR03


IsoR03 Profile in W vs. Z sample

- Isolation profile in Pt and Eta ($iso < 2.0$) - IsoR03 : W vs. Z

Isolation E (R=0.3) vs Muon Pt



Isolation E (R=0.3) vs Muon η



- Not much difference b/w W and Z sample
 - will check isoE distribution itself, not profile plot

Summary

- IsolationE in Pt and Eta
 - Pt dependence is small
 - IsoR03 has a large eta dependence in the endcap region
 - IsoR03fw is relatively flat compared to IsoR03
 - IsoR03fw is better to use for eta dependence
 - efficiency factor will be canceled out in lepton charge asymmetry
 - isolation shape to measure the background has a issue with isoR03
- Iso shape does not have much difference between W and Z sample
- Need to decide what Isolation variable to use for the measurement